



SELECTING PLANTS FOR POLLINATORS



A REGIONAL GUIDE FOR FARMERS, LAND MANAGERS, AND GARDENERS IN THE



OUTER COASTAL PLAIN MIXED PROVINCE

INCLUDING THE STATES OF:
DELAWARE, FLORIDA, GEORGIA,
LOUISIANA, NORTH CAROLINA,
SOUTH CAROLINA

AND PARTS OF
ALABAMA, MARYLAND,
MISSISSIPPI, TEXAS,
AND VIRGINIA



**POLLINATOR
PARTNERSHIP**

Protect their lives. Preserve ours.

TABLE OF CONTENTS

| | |
|---|-----------|
| WHY SUPPORT POLLINATORS? | 4 |
| GETTING STARTED | 5 |
| OUTER COASTAL PLAIN MIXED PROVINCE | 6 |
| MEET THE POLLINATORS | 8 |
| PLANT TRAITS | 10 |
| DEVELOPING PLANTINGS | 12 |
| FARMS | 13 |
| PUBLIC LANDS | 14 |
| HOME LANDSCAPES | 15 |
| BLOOM PERIODS | 16 |
| PLANTS THAT ATTRACT POLLINATORS | 18 |
| HABITAT HINTS | 20 |
| CHECKLIST | 22 |
| RESOURCES AND FEEDBACK | 23 |

This is one of several guides for different regions in the United States. We welcome your feedback to assist us in making the future guides useful. Please contact us at feedback@pollinator.org

SELECTING PLANTS FOR POLLINATORS

A REGIONAL GUIDE FOR FARMERS, LAND MANAGERS, AND GARDENERS

IN THE ECOLOGICAL REGION OF THE **OUTER COASTAL PLAIN MIXED PROVINCE**

INCLUDING THE STATES OF:

DELAWARE, FLORIDA, GEORGIA, LOUISIANA,
NORTH CAROLINA, SOUTH CAROLINA

AND PARTS OF

ALABAMA, MARYLAND, MISSISSIPPI, TEXAS,
AND VIRGINIA

A NAPPC AND POLLINATOR PARTNERSHIP PUBLICATION

This guide was funded by the National Fish and Wildlife Foundation, the C.S. Fund, the Plant Conservation Alliance, the U.S. Forest Service, and the Bureau of Land Management with oversight by Pollinator Partnership (pollinator.org), in support of the North American Pollinator Protection Campaign (NAPPC-napcc.org).



WHY SUPPORT POLLINATORS?

IN THEIR 1996 BOOK, *THE FORGOTTEN POLLINATORS*, Buchmann and Nabhan estimated that animal pollinators are needed for the reproduction of 90% of flowering plants and one third of human food crops. Each of us depends on these industrious pollinators in a practical way to provide us with the wide range of foods we eat. In addition, pollinators are part of the intricate web that supports the biological diversity in natural ecosystems that helps sustain our quality of life.

Abundant and healthy populations of pollinators can improve fruit set and quality, and increase fruit size. In farming situations this increases production per acre. In the wild, biodiversity increases and wildlife food sources increase.

Apples, tomatoes, citrus, melons, and strawberries are some of the crops raised in the Outer Coastal Plain Mixed Province that rely on honey bees and native bees for pollination. Domestic honey bees pollinate approximately \$10 billion worth of crops in the U.S. each year.

Unfortunately, the numbers of both native pollinators and domesticated bee populations are declining. They are threatened by habitat loss, disease, and the excessive and inappropriate use of pesticides. The loss of commercial bees to Colony Collapse Disorder (CCD) has highlighted how severe the issues of proper hive management are to reduce stresses caused by disease, pesticide use, insufficient nutrition, and transportation practices. Currently, the pollination services that the commercial beekeeping industry provides are receiving much needed research and conservation resources. The efforts to understand the threats to commercial bees should help us understand other pollinators and their roles in the environment as well.

It is imperative that we take immediate steps to help pollinator populations thrive. The beauty of the situation is that by supporting pollinators' need for habitat, we support our own needs for food and support diversity in the natural world.

Thank you for taking time to consult this guide. By adding plants to your landscape that provide food and shelter for pollinators throughout their active seasons and by adopting pollinator friendly landscape practices, you can make a difference to both the pollinators and the people that rely on them.

Kelly Bills
Executive Director
Pollinator Partnership

**“ FARMING FEEDS
THE WORLD, AND
WE MUST REMEMBER
THAT POLLINATORS
ARE A CRITICAL
LINK IN OUR FOOD
SYSTEMS.”**

-- PAUL GROWALD,
CO-FOUNDER,
POLLINATOR PARTNERSHIP



GETTING STARTED



THIS REGIONAL GUIDE IS just one in a series of plant selection tools designed to provide information on how individuals can influence pollinator populations through choices they make when they farm a plot of ground, manage large tracts of public land, or plant a garden. Each of us can have a positive impact by providing the essential habitat requirements for pollinators including food, water, shelter, and enough space to allow pollinators to raise their young.

Pollinators travel through the landscape without regard to property ownership or state boundaries. We've chosen to use R.G. Bailey's classification system to identify the geographic focus of this guide and to underscore the connections between climate and vegetation types that affect the diversity of pollinators in the environment.

Bailey's Ecoregions of the United States, developed by the United States Forest Service, is a system

created as a management tool and is used to predict responses to land management practices throughout large areas. This guide addresses pollinator-friendly land management practices in what is known as the Outer Coastal Plain Mixed Province.

Portions of eleven states make up the 173,800 square miles of this generally flat province with elevations ranging from sea level to around 300 feet. The topography is primarily gently sloping and made up of many slowly moving streams, lakes, marshes, and swamps. Average annual temperatures are mild, ranging from 60° to 70°F.

This moist, green province is characterized by rainfall throughout the year, between 40 to 60 inches annually, creating in many areas what is known as a temperate rainforest. This type of rainforest has fewer species of vegetation than its tropical counterpart and therefore has larger populations of individual species. Common

species include evergreen oaks and members of the laurel and magnolia families. Tree trunks are often covered in moss at higher elevations, and a well-developed lower layer of vegetation, including tree ferns, shrubs, and herbaceous plants is often present.

Long before there were homes and farms in this area, the original, natural vegetation provided continuous cover and adjacent feeding opportunities for wildlife, including pollinators. In choosing plants, aim to create habitat for pollinators that allow adequate food shelter, and water sources. Most pollinators have very small home ranges. You can make a difference by understanding the vegetation patterns of the farm, forest, or neighbor's yard adjacent to you and by making planting choices that support the pollinators' need for food and shelter as they move through the landscape.


UNDERSTANDING THE OUTER COASTAL PLAIN MIXED PROVINCE



- ✿ This region is designated **number 232** in the Baileys' Ecosystem Provinces. To see a map of the provinces go to: fs.fed.us/colorimagemap/ecoreg1_provinces.html
- ✿ Not sure about which bioregion you live or work in? Go to pollinator.org and click on **Ecoregion Locator** for help.
- ✿ 173,800 square miles within 11 states.
- ✿ Primarily flat or gently sloping and characterized by marshes, swamps, lakes, and slow-moving streams.
- ✿ Elevations ranging from sea level to around 300 feet.
- ✿ Average annual temperature range from 60° to 70°F.
- ✿ Average year-round precipitation between 40-60 inches.
- ✿ USDA Hardiness Zones 7a-9b (1990 version).

CHARACTERISTICS

- ✿ Dominated by evergreen oaks, laurels, magnolias.
- ✿ Usually a well-developed lower layer of growth including tree ferns, small palms, shrubs, and herbaceous plants, including many lianas and epiphytes.
- ✿ Trees are often covered in moss at higher elevations where there are clouds and fog.
- ✿ Gum and cypress are abundant along the Atlantic Coast.
- ✿ Subclimax pine forests dominate upland areas, and usually have an understory of grasses and sedges called savannas.



The Outer Coastal Plain
Mixed Province

includes the states of:

Delaware, Florida, Georgia,
Louisiana, North Carolina,
South Carolina

And parts of:

Alabama, Maryland, Mississippi,
Texas, and Virginia

**“ ADDING NATIVE PLANTINGS IN RIPARIAN AREAS
TO IMPROVE POLLINATOR HABITAT MAKES
SENSE IN ADVANCING OUR FAMILY FARM’S
CONSERVATION AND ECONOMIC OBJECTIVES,
ENHANCING BENEFICIAL WILDLIFE AND
IMPROVING POLLINATION IN OUR ORCHARD
AND GARDEN. ”**

—LEE MCDANIEL, FARMER AND PRESIDENT,
NATIONAL ASSOCIATION OF CONSERVATION DISTRICTS

MEET THE POLLINATORS



Photos courtesy www.dangphoto.net

Tiger Swallowtail butterfly on Purple Hollyhock, (above), and a honey bee on a large holly bush in the Atlanta, Georgia area.



WHO ARE THE POLLINATORS?

BEES

Bees are well documented pollinators in the natural and agricultural systems of the Outer Coastal Plain Mixed Province and the Inner Coastal Plain. A wide range of crops including apples, citrus, melons, strawberries, and tomatoes are just a few plants that benefit from bee pollinators.

Most of us are familiar with the colonies of honey bees that have been the workhorses of agricultural pollination for years in the United States. They were imported from Europe almost 400 years ago.

There are nearly 4000 species of native ground and twig nesting bees in the U.S. Some form colonies while others live and work a solitary life. Native bees currently pollinate many crops and can be encouraged to do more to support agricultural endeavors if their needs for nesting habitat are met and if suitable sources of nectar, pollen, and water are provided. Bees have tongues of varying lengths that help determine which flowers they can obtain nectar and pollen from.

The bumble bee (*Bombus* spp.) forms small colonies, usually underground. They are generalists, feeding on a wide range of plant material from February to November and are important pollinators of tomatoes. The sweat bee (family *Halictidae*) nests


underground. Various species are solitary while others form loose colonies.

Solitary bees include carpenter bees (*Xylocopa* spp.), which nest in wood; digger, or polyester bees (*Colletes* spp.), which nest underground; leafcutter bees (*Megachile* spp.), which prefer dead trees or branches for their nest sites; and mason bees (*Osmia* spp.), which utilize cavities that they find in stems and dead wood. Cactus bees (*Diadasia* spp.) are also solitary ground nesters.

BUTTERFLIES

Gardeners have been attracting butterflies to their gardens for some time. These insects tend to be eye-catching, as are the flowers that attract them. Position flowering plants where they have full sun and are protected from the wind. Also, you will need to provide open areas (e.g. bare earth, large stones) where butterflies may bask, and moist soil from which they may get needed minerals. By providing a safe place to eat and nest, gardeners can also support the pollination role that butterflies play in the landscape. It might mean accepting slight damage to the plants, known as host plants, that provide food for the larval stage of the butterfly.

A diverse group of butterflies are present in garden areas and woodland edges that provide bright flowers, water sources, and specific host plants. Numerous trees, shrubs, and herbaceous plants support butterfly populations.



Butterflies are in the Order *Lepidoptera*. Some of the species in the Outer Coastal Plain Mixed Province are Brush-footed, Gossamer-winged, Swallowtail, Parnassian, Skipper, White, Sulphur and Milkweed butterflies. They usually look for flowers that provide a good landing platform.

Wet mud areas provide butterflies with both the moisture and minerals they need to stay healthy. Butterflies eat rotten fruit and even dung, so don't clean up all the messes in your garden!

MOTHS

Moths are most easily distinguished from butterflies by their antennae. Butterfly antennae are simple with a swelling at the end. Moth antennae differ from simple to featherlike, but never have a swelling at the tip. In addition, butterflies typically are active during the day; moths at night. Butterfly bodies are not very hairy, while moth bodies are quite hairy and more stout.

Moths, generally less colorful than butterflies, also play a role in pollination. They are attracted to flowers that are strongly sweet smelling, open in late afternoon or night, and are typically white or pale colored.

BEETLES

Over 30,000 species of beetles are found in the United States and many of them can be found on flower heads. Gardeners have yet to intentionally draw beetles

to their gardens, possibly because beetle watching isn't as inspiring as butterfly or bird watching. Yet beetles do play a role in pollination. Some have a bad reputation because they can leave a mess behind, damaging plant parts that they eat. Beetles are not as efficient as some pollinators. They wander between different species, often dropping pollen as they go. Beetle pollinated plants tend to be large, strong scented flowers with their sexual organs exposed. They are known to pollinate Magnolia, sweetshrub (*Calycanthus*), paw paws, and yellow pond lilies.

FLIES

It may be hard to imagine why one would want to attract flies to the garden. However, like beetles, the number of fly species and the fact that flies are generalist pollinators (visit many species of plants), should encourage us all to leave those flies alone and let them do their job as pollinators.

Recent research indicates that flies primarily pollinate small flowers that bloom under shade and in seasonally moist habitats. The National Research Council's *Status of Pollinators in North America* study states that flies are economically important as pollinators for a range of annual and bulbous ornamental flowers.

Plants pollinated by the fly include the American pawpaw (*Asimina triloba*), dead horse arum (*Helicodiceros muscivorus*), skunk cabbage (*Symplocarpus foetidus*),

goldenrod (*Solidago* spp.), and members of the carrot family like Queen Anne's lace (*Daucus carota*).

BIRDS

Hummingbirds are the primary birds which play a role in pollination in North America. Their long beaks and tongues draw nectar from tubular flowers. Pollen is carried on both the beaks and feathers of different hummingbirds. The regions closer to the tropics, with warmer climates, boast the largest number of hummingbird species and the greatest number of native plants to support the bird's need for food. White-winged doves (*Zenaida asiatica*) are also pollinators of the saguaro cactus (*Carnegeia gigantea*) in the south central United States.

Bright colored tubular flowers attract hummingbirds to gardens throughout the United States. Hummingbirds can see the color red; bees cannot. Many tropical flowers, grown as perennials in the Outer Coastal Plain Mixed Province, along with native woodland edge plants, attract hummingbirds.

BATS

Though bats in the Outer Coastal Plain Mixed Province are not pollinators, bats play an important role in pollination in the southwest where they feed on agave and cactus. The long-nosed bats' head shape and long tongue allows it to delve into flower blossoms and extract both pollen and nectar.



PLANT TRAITS

WHICH FLOWERS DO THE POLLINATORS PREFER?

NOT ALL POLLINATORS ARE found in each North American province, and some are more important in different parts of the United States. Use this page as a resource to understand the plants and pollinators where you live.

Plants can be grouped together based on the similar characteristics of their flowers. These floral characteristics can be useful to predict the type of pollination method or animal that is most effective for that group of plants. This association between floral characteristics and pollination method is called a pollination syndrome.

The interactions of animal pollinators and plants have influenced the evolution of both groups of organisms. A mutualistic relationship between the pollinator and the plant species helps the pollinator find necessary pollen and nectar sources and helps the plant reproduce by ensuring that pollen is carried from one flower to another.

| Plant Trait | Bats | Bees | Beetles |
|---------------|--|---|----------------------------------|
| Color | Dull white, green or purple | Bright white, yellow, blue, or UV | Dull white or green |
| Nectar guides | Absent | Present | Absent |
| Odor | Strong musty; emitted at night | Fresh, mild, pleasant | None to strongly fruity or fetid |
| Nectar | Abundant; somewhat hidden | Usually present | Sometimes present; not hidden |
| Pollen | Ample | Limited; often sticky and scented | Ample |
| Flower Shape | Regular; bowl shaped – closed during day | Shallow; have landing platform; tubular | Large bowl-like, Magnolia |

This chart and more information on pollinator syndromes can be found at:



AND THE POLLINATORS THEY ATTRACT

Pollinator

| Birds | Butterflies | Flies | Moths | Wind |
|---|---|---|--|---|
| Scarlet, orange, red or white | Bright, including red and purple | Pale and dull to dark brown or purple; flecked with translucent patches | Pale and dull red, purple, pink or white | Dull green, brown, or colorless; petals absent or reduced |
| Absent | Present | Absent | Absent | Absent |
| None | Faint but fresh | Putrid | Strong sweet; emitted at night | None |
| Ample; deeply hidden | Ample; deeply hidden | Usually absent | Ample; deeply hidden | None |
| Modest | Limited | Modest in amount | Limited | Abundant; small, smooth, and not sticky |
| Large funnel like; cups, strong perch support | Narrow tube with spur; wide landing pad | Shallow; funnel like or complex and trap-like | Regular; tubular without a lip | Regular; small and stigmas exerted |

<http://fs.fed.us/wildflowers/pollinators/syndromes.shtml>



DEVELOPING LANDSCAPE PLANTINGS THAT PROVIDE POLLINATOR HABITAT

WHETHER YOU ARE A FARMER of many acres, land manager of a large tract of land, or a gardener with a small lot, you can increase the number of pollinators in your area by making conscious choices to include plants that provide essential habitat for bees, butterflies, moths, beetles, hummingbirds and other pollinators.

FOOD:

Flowers provide nectar (high in sugar and necessary amino acids) and pollen (high in protein) to pollinators.

Fermenting fallen fruits also provide food for bees, beetles and butterflies.

Specific plants, known as host plants, are eaten by the larvae of pollinators such as butterflies.

- Plant in groups to increase pollination efficiency. If a pollinator can visit the same type of flower over and over, it doesn't have to relearn how to enter the flower and can transfer pollen to the same species, instead of squandering the pollen on unreceptive flowers.
- Plant with bloom season in mind, providing food from early spring to late fall. (see Bloom Periods pp. 16-17)
- Plant a diversity of plants to support a variety of pollinators. Flowers of different color, fragrance, and season of bloom on plants of different heights will attract different pollinator species and provide pollen and nectar throughout the seasons.
- Many herbs and annuals, although

not native, are very good for pollinators. Mint, oregano, garlic, chives, parsley and lavender are just a few herbs that can be planted. Old fashioned zinnias, cosmos, and single sunflowers support bees and butterflies.

- Recognize weeds that might be a good source of food. For example, dandelions provide nectar in the early spring before other flowers open. Plantain is alternate host for the Baltimore Checkerspot.
- Learn and utilize Integrated Pest Management (IPM) practices to address pest concerns. Minimize or eliminate the use of pesticides.

SHELTER:

Pollinators need protection from severe weather and from predators as well as sites for nesting and roosting.

- Incorporate different canopy layers in the landscape by planting trees, shrubs, and different-sized perennial plants.
- Leave dead snags for nesting sites of bees, and other dead plants and leaf litter for shelter.
- Build bee boxes to encourage solitary, non-aggressive bees to nest on your property.
- Leave some areas of soil uncovered to provide ground nesting insects easy access to underground tunnels.
- Group plantings so that pollinators can move safely through the landscape protected from predators.
- Include plants that are needed

by butterflies during their larval development.

WATER:

A clean, reliable source of water is essential to pollinators.

- Natural and human-made water features such as running water, pools, ponds, and small containers of water provide drinking and bathing opportunities for pollinators.
- Ensure the water sources have a shallow or sloping side so the pollinators can easily approach the water without drowning.

Your current landscape probably includes many of these elements. Observe wildlife activity in your farm fields, woodlands, and gardens to determine what actions you can take to encourage other pollinators to feed and nest. Evaluate the placement of individual plants and water sources and use your knowledge of specific pollinator needs to guide your choice and placement of additional plants and other habitat elements. Minor changes by many individuals can positively impact the pollinator populations in your area. Watch for - and enjoy - the changes in your landscape!

- **CAUTION:** Remember that pesticides are largely toxic to pollinators. Extreme caution is warranted if you choose to use any pesticide. Strategically apply pesticides only for problematic target species.



FARMS

Tomatoes, citrus, melons, apples and strawberries are a few of the food crops in the Outer Coastal Plain Mixed Province that will benefit from strong native bee populations that boost pollination efficiency. Incorporate different plants throughout the farm that provide food for native populations when targeted crops are not in flower.

Farmers have many opportunities to incorporate pollinator-friendly land management practices on their land which will benefit the farmer in achieving his or her production goals:

- Manage the use of pesticides to reduce the impact on native pollinators. Spray when bees aren't active (just after dawn) and choose targeted ingredients.
- Carefully consider the use of herbicides. Perhaps the targeted

weeds can provide needed food for pollinators.

- Minimize tillage to protect ground nesting pollinators.
- Ensure water sources are scattered throughout the landscape.
- Choose a variety of native plants to act as windbreaks, riparian buffers, and field borders throughout the farm.
- Plant unused areas of the farm with temporary cover crops that can provide food or with a variety of trees, shrubs, and flowers that provide both food and shelter for pollinators.
- Check with your local Natural Resources Conservation Service (NRCS) office to see what technical and financial support might be available to assist you in your effort to provide nectar, pollen, and larval food sources for pollinators on your farm.

“**FOOD SUPPLIES FOR BEES ARE CRITICAL TO MAINTAINING STRONG HIVES FOR ALMOND POLLINATION THE FOLLOWING WINTER.**”

-- DAN CUMMINGS,
CHICO, CALIFORNIA
ALMOND GROWER.



Illustrations by Carolyn Vibbert

PUBLIC LANDS

“FROM HUMMINGBIRDS TO BEETLES, TO BUTTERFLIES, NATURE’S POLLINATORS HELP KEEP MIDWIN’S TALLGRASS PRAIRIE RESTORATIONS FULL OF DIVERSE FLOWERING PLANTS. INSECT MONITORING PROVIDES A KEY MEASURE OF OUR SUCCESS.”

-- LOGAN LEE
PRAIRIE SUPERVISOR, MIDWIN
NATIONAL TALLGRASS PRAIRIE



Public lands are maintained for specific reasons ranging from high impact recreation to conservation. In the Outer Coastal Plain Mixed Province, natural areas have been altered to allow for roads, buildings, open lawn areas, boat ramps, and vistas. Less disturbed natural areas can be augmented with plantings of native plant species. Existing plantings around buildings and parking areas should be evaluated to determine if pollinator-friendly plants can be substituted or added to attract and support pollinators. Public land managers have a unique opportunity to use their plantings as an education tool to help others understand the importance of pollinators in the environment through signs, brochures, and public programs.

In an effort to increase populations of pollinators the land manager can:

- Inventory and become knowledgeable of local pollinators.
- Provide connectivity between vegetation areas by creating corridors of perennials, shrubs, and trees that provide pollinators shelter and food as they move through the landscape.
- Maintain a minimum of lawn areas that support recreational needs.
- Restrict the use of pesticides and herbicides.
- Provide water sources in large open areas.
- Maintain natural meadows and openings that provide habitats for sun-loving wildflowers and grasses.
- Remove invasive species and encroaching shrubs and trees.



HOME LANDSCAPES

“ A GARDEN IS ONLY AS RICH AND BEAUTIFUL AS THE INTEGRAL HEALTH OF THE SYSTEM; POLLINATORS ARE ESSENTIAL TO THE SYSTEM - MAKE YOUR HOME THEIR HOME. ”

-- DERRY MACBRIDE
NATIONAL AFFAIRS AND
LEGISLATION CHAIRWOMAN,
GARDEN CLUB OF AMERICA

Gardeners have a wide array of plants to use in their gardens. Native plants, plants introduced from years of plant exploration from around the world, and plants developed by professional and amateur breeders can be found in garden centers, in catalogs, and on web-sites. Use your knowledge of pollinator needs to guide your choices.

- Choose a variety of plants that will provide nectar and pollen throughout the growing season.
- Resist the urge to have a totally manicured lawn and garden. Leave bare ground for ground nesting bees. Leave areas of dead wood and leaf litter for other insects.
- Strive to eliminate the use of all pesticides.
- Find local resources to help you in your efforts. Contact your local county extension agent or native plant society. Visit your regional botanic gardens and arboreta.

The scale of your plantings will vary but it is important to remember that you are trying to provide connectivity to the landscape adjacent to your property. Don't just look within your property boundaries. If your neighbor's property provides an essential element, such as water, which can be utilized by pollinators visiting your land, you may be able to devote more space to habitat elements that are missing nearby. It is best to use native plants which have evolved to support the needs of specific native pollinators. Some pollinators, however, are generalists and visit many different plants, both native and non-native. Be sure that any non-native plants you choose to use are not invasive. Remember that specialized cultivars sometimes aren't used by pollinators. Flowers that have been drastically altered, such as those that are double or a completely different color than the wild species, often prevent pollinators from finding and feeding on the flowers. In addition, some altered plants don't contain the same nectar and pollen resources that attract pollinators to the wild types.

- **CAUTION:** Take time to evaluate the source of your plant material. You want to ensure you get plants that are healthy and correctly identified. Your local native plant society can help you make informed decisions when searching for plants.



BLOOM PERIODS

FOR THE OUTER COASTAL PLAIN MIXED PROVINCE

The following chart lists plants and the time they are in bloom throughout the growing seasons. Choose a variety of flower colors and make sure something is blooming at all times! Note for all charts: When more than one species of the same genus is useful, the genus name is followed by “spp.”

| Botanical Name | Common Name | | | March | April | May | June | July | Aug | Sept | Oct | Nov |
|---|----------------------|---------------------|-------------|---------------|---------------------|---------------------|----------------|----------------|--------------|------------|-------|-----|
| Trees & Shrubs | | | | | | | | | | | | |
| <i>Agarista populifolia</i> | Florida Leucothoe | white | May-June | | | white | white | | | | | |
| <i>Aesculus parviflora</i> | Bottlebrush Buckeye | white | July-August | | | | | white | white | | | |
| <i>Aesculus pavia</i> | Red Buckeye | red, reddish-yellow | April-May | | red, reddish-yellow | red, reddish-yellow | | | | | | |
| <i>Aronia arbutifolia</i> | Red Chokeberry | white | March-May | white | white | white | | | | | | |
| <i>Asimina parviflora</i> | Small-fruited Pawpaw | reddish-maroon | April-May | | reddish-maroon | reddish-maroon | | | | | | |
| <i>Baccharis halimifolia</i> | Groundsel Bush | white | Sept-Oct | | | | | | | white | white | |
| <i>Callicarpa americana</i> | American Beautyberry | lavender, pink | June-July | | | | lavender, pink | lavender, pink | | | | |
| <i>Clethra alnifolia</i> | Sweet Pepperbush | white | July | | | | | white | | | | |
| <i>Cliftonia monophylla</i> | Black Ti-ti | white or pink | March-April | white or pink | white or pink | | | | | | | |
| <i>Clinopodium georgianum</i> | Georgia Savory | light pink | July-Sept | | | | | light pink | light pink | light pink | | |
| <i>Clinopodium coccineum</i> | Red Savory | red | July-Sept | | | | | red | red | red | | |
| <i>Crataegus aestivalis</i> | Mayhaw | white | March-April | white | white | | | | | | | |
| <i>Cyrilla racemiflora</i> | Ti-ti, Leatherwood | white | May-July | | | white | white | white | | | | |
| <i>Erythrina herbacea</i> | Coral Bean | red | May-July | | | red | red | red | | | | |
| <i>Eubotrys racemosa</i> | Coastal Fetterbush | white | April-May | | white | white | | | | | | |
| <i>Fothergilla gardenii</i> | Fothergilla | white | March-May | white | white | white | | | | | | |
| <i>Gordonia lasianthus</i> | Loblolly Bay | white | July-Sept | | | | | white | white | white | | |
| <i>Halesia diptera</i> var. <i>diptera</i> | Two Wing Silverbell | white | April-May | | white | white | | | | | | |
| <i>Ilex coriacea</i> | Big Gallberry | white | March-May | white | white | white | | | | | | |
| <i>Ilex decidua</i> var. <i>decidua</i> | Possum-haw | white | March-May | white | white | white | | | | | | |
| <i>Ilex glabra</i> | Inkberry | white | May-June | | | white | white | | | | | |
| <i>Ilex vomitoria</i> | Yaupon | white | March-May | white | white | white | | | | | | |
| <i>Illicium floridanum</i> | Purple Anise | maroon | April-May | | maroon | maroon | | | | | | |
| <i>Itea virginica</i> | Virginia Sweetspire | white | May-June | | | white | white | | | | | |
| <i>Lyonia lucida</i> | Shining Fetterbush | white | April-May | | white | white | | | | | | |
| <i>Magnolia grandiflora</i> | Southern Magnolia | white | May-June | | | white | white | | | | | |
| <i>Magnolia virginiana</i> var. <i>virginiana</i> | Sweet Bay | white | April | | white | | | | | | | |
| <i>Morella cerifera</i> | Common Wax Myrtle | white | April | | white | | | | | | | |
| <i>Rhododendron alabamense</i> | Alabama Azalea | white | March-April | white | white | | | | | | | |
| <i>Rhododendron atlanticum</i> | Coastal Azalea | pinkish white | April-May | | pinkish white | pinkish white | | | | | | |
| <i>Rhododendron austrinum</i> | Florida Azalea | yellowish-orange | April | | yellowish-orange | | | | | | | |
| <i>Persea borbonia</i> | Red Bay | yellow, small | May-June | | | yellow, small | yellow, small | | | | | |
| <i>Pinckneya bracteata</i> | Fever-tree | creamy rose | May-June | | | creamy rose | creamy rose | | | | | |
| <i>Sabal minor</i> | Dwarf Palmetto | white | June-July | | | | white | white | | | | |
| <i>Sesbania drummondii</i> | Poisonbean | light yellow | August | | | | | | light yellow | | | |
| <i>Stewartia malacodendron</i> | Silky Camellia | white | May-June | | | white | white | | | | | |
| <i>Styrax americanus</i> var. <i>americanus</i> | American Snowbell | white | April-June | | white | white | white | | | | | |
| <i>Styrax grandifolius</i> | Bigleaf Snowbell | white | April-May | | white | white | | | | | | |
| <i>Vaccinium crassifolium</i> | Creeping Blueberry | white | April-May | | white | white | | | | | | |
| <i>Viburnum nudum</i> | Possumhaw | white | April-May | | white | white | | | | | | |

| Botanical Name | Common Name | | | March | April | May | June | July | Aug | Sept | Oct | Nov |
|--|---------------------------|--------------------------------|-------------|-------------------------------|-------------------------------|-------------------------------|---------------|--------------------------------|--------------------------------|---------------------------|---------------------------|--------|
| <i>Zenobia pulverulenta</i> | Zenobia | white | April-June | | white | white | white | | | | | |
| Perennial Flowers | | | | | | | | | | | | |
| <i>Asclepias lanceolata</i> | Red Milkweed | orange-red | June-August | | | | | | yellow | yellow | yellow | |
| <i>Bigelovia nudata</i> | Rayless Goldenrod | yellow | Aug-Oct | | | | | | white/pink | white/pink | white/pink | |
| <i>Boltonia caroliniana</i> | Carolina Doll's-daisy | white/pink | Aug-Oct | | | | | purple to pink | purple to pink | purple to pink | purple to pink | |
| <i>Carphephorus odoratissimus</i> | Vanilla Plant | purple to pink | July-Oct | | | | | | | | | |
| <i>Cirsium horridulum</i> | Yellow Thistle | pale yellow or purple | March-May | pale yellow or purple | pale yellow or purple | pale yellow or purple | | | | | | |
| <i>Conradina canescens</i> | Gray Rosemary | light purple | January-May | light purple | light purple | light purple | | | | | | |
| <i>Chrysogonum virginianum</i> var. <i>australe</i> | Gulf Coast Green-and-gold | yellow | March-May | yellow | yellow | yellow | | | | | | |
| <i>Dyschoriste oblongifolia</i> | Blue Twin Flower | blue | April-May | | blue | blue | | | | | | |
| <i>Coreopsis falcata</i> | Pool Coreopsis | yellow | May-July | | | yellow | yellow | yellow | | | | |
| <i>Eryngium aquaticum</i> | Marsh Eryngo | purplish-blue | July-Sept | | | | | purplish-blue | purplish-blue | purplish-blue | | |
| <i>Eurybia spectabilis</i> | Showy Aster | purple with yellow center | Aug-Oct | | | | | | purple with yellow center | purple with yellow center | purple with yellow center | |
| <i>Helenium pinnatifidum</i> | Savanna Sneezeweed | yellow | April-May | | yellow | yellow | | | | | | |
| <i>Helianthus angustifolius</i> | Narrowleaf Sunflower | yellow | Sept-Oct | | | | | | | yellow | yellow | |
| <i>Helianthus heterophyllus</i> | Savanna Sunflower | yellow | Aug-Oct | | | | | | yellow | yellow | yellow | |
| <i>Hibiscus coccineus</i> | Scarlet Hibiscus | red | July-Aug | | | | | red | red | | | |
| <i>Hibiscus grandiflorus</i> | Swamp Rose-mallow | pink/white | July-Aug | | | | | pink/white | pink/white | | | |
| <i>Hibiscus moscheutos</i> | Eastern Rose-mallow | white to pink with red throats | June-Aug | | | | | white to pink with red throats | white to pink with red throats | | | |
| <i>Iris verna</i> var. <i>verna</i> | Coastal Plain Dwarf Iris | blue with golden yellow sepal | March-May | blue with golden yellow sepal | blue with golden yellow sepal | blue with golden yellow sepal | | | | | | |
| <i>Iris virginica</i> | Southern Blueflag | blue | April-May | | blue | blue | | | | | | |
| <i>Kosteletskyia virginica</i> var. <i>virginica</i> | Marsh Mallow | pink | July-Oct | | | | | pink | pink | pink | pink | |
| <i>Liatris elegans</i> | Elegant Blazing-star | rosy-pink | Sept-Oct | | | | | | | rosy-pink | rosy-pink | |
| <i>Lobelia elongata</i> | Blue Lobelia | blue | Aug-Oct | | | | | | blue | blue | blue | |
| <i>Monarda punctata</i> var. <i>punctata</i> | Eastern Horse-mint | yellow/purplish-pink | Aug-Sept | | | | | | yellow/purplish-pink | yellow/purplish-pink | | |
| <i>Orontium aquaticum</i> | Golden Club | yellow | March-April | yellow | yellow | | | | | | | |
| <i>Pontederia cordata</i> var. <i>cordata</i> | Pickerelweed | purple | June-Aug | | | | purple | purple | purple | | | |
| <i>Pityopsis graminifolia</i> var. <i>tenuifolia</i> | Narrow-leaved Silkgrass | yellow | Sept-Oct | | | | | | | yellow | yellow | |
| <i>Rhexia alifanus</i> | Smooth Meadow-beauty | pink | May-Sept | | | pink | pink | pink | pink | pink | | |
| <i>Solidago sempivirens</i> var. <i>mexicana</i> | Seaside Goldenrod | yellow | Sept-Nov | | | | | | | yellow | yellow | yellow |
| <i>Stokesia laevis</i> | Stoke's Aster | blue | July-Aug | | | | | blue | blue | | | |
| <i>Yucca filamentosa</i> | Spoonleaf Yucca | white/green | April-June | | white/green | white/green | white/green | | | | | |
| <i>Zephyranthes atamasca</i> | Atamasco Lily | white | March-April | white | white | | | | | | | |
| Vines | | | | | | | | | | | | |
| <i>Ampelaster carolinianus</i> | Climbing Aster | rose-pink | Sept-Oct | | | | | | | rose-pink | rose-pink | |
| <i>Bignonia capreolata</i> | Cross-vine | red and yellow | April-May | | red and yellow | red and yellow | | | | | | |
| <i>Campsis radicans</i> | Trumpet-creeper | orange | June-July | | | | orange | orange | | | | |
| <i>Clematis crispa</i> | Marsh Clematis | bluish-purple | June-August | | | | bluish-purple | bluish-purple | bluish-purple | | | |
| <i>Wisteria frutescens</i> | American Wisteria | bluish-purple, lavender | April-May | | bluish-purple, lavender | bluish-purple, lavender | | | | | | |
| <i>Gelsemium rankinii</i> | Swamp Jessamine | yellow | March-April | yellow | yellow | | | | | | | |
| <i>Ipomoea sagittata</i> | Arrow-leaf Morning Glory | pink | July-Sept | | | | | pink | pink | pink | | |

PLANTS THAT ATTRACT POLLINATORS IN THE OUTER COASTAL PLAIN MIXED PROVINCE

The following chart lists plants that attract pollinators. It is not exhaustive, but provides guidance on where to start. Annuals, herbs, weeds, and cover crops provide food and shelter for pollinators, too.

| Botanical Name | Common Name | Color | Height | Flower Season | Sun | Soil | Visitation by pollinator | Also a host |
|---|----------------------|---------------------|--------|---------------|------------------------|-----------------------------|--|--|
| Trees & Shrubs | | | | | | | | |
| <i>Aesculus parviflora</i> | Bottlebrush Buckeye | white | 6-12' | July-August | shade to partial shade | moist, well drained | butterflies (zebra swallowtail), bumblebees | |
| <i>Aesculus pavia</i> | Red Buckeye | red, reddish-yellow | 20-25' | April-May | shade to partial shade | moist | hummingbirds, bees, bumblebees | |
| <i>Aronia arbutifolia</i> | Red Chokeberry | white | to 15' | March-May | sun to partial shade | moist to wet | butterflies, bees | butterflies (Coral Hairstreak, Striped Hairstreak), moths (Bluish Spring, Praeclara Underwing) |
| <i>Asimina parviflora</i> | Small-fruited Pawpaw | reddish-maroon | 6-9' | April-May | shade to partial shade | dry to moist | flies (fruit flies, green bottle flies), beetles, moths | |
| <i>Callicarpa americana</i> | American Beautyberry | lavender, pink | 4-6' | June-July | sun to partial shade | dry to moist | bees, butterflies | |
| <i>Clethra alnifolia</i> | Sweet Pepperbush | white | 4-9' | July | sun | moist to wet, acidic | butterflies (Great Purple Hairstreak, Snout butterfly), bees (honeybees, others), hummingbirds | |
| <i>Clinopodium georgianum</i> | Georgia Savory | light pink | 12-18" | July-Sept | sun to partial shade | dry | bees | |
| <i>Crataegus aestivalis</i> | Mayhaw | white | 12-18' | March-April | shade to partial shade | mesic to wet | flies (midges, others) | butterflies |
| <i>Cyrilla racemiflora</i> | Ti-ti, Leatherwood | white | 5-15' | May-July | sun to partial shade | moist to wet | bees (honeybees, others), butterflies | |
| <i>Erythrina herbacea</i> | Coral Bean | red | to 24' | May-July | sun to partial shade | dry | butterflies | |
| <i>Fothergilla gardenii</i> | Fothergilla | white | 2-4' | March-May | sun to partial shade | moist | bees (honeybees, others) | |
| <i>Gordonia lasianthus</i> | Loblolly Bay | white | to 75' | July-Sept | sun to shade | moist to wet, acidic | bees | |
| <i>Halesia diptera</i> var. <i>diptera</i> | Two Wing Silverbell | white | to 30' | April-May | sun to partial shade | moist to wet | bees, butterflies | |
| <i>Ilex coriacea</i> | Big Gallberry | white | to 15' | March-May | sun to partial shade | wet | bees | butterflies |
| <i>Ilex decidua</i> var. <i>decidua</i> | Possum-haw | white | to 20' | March-May | shade to partial shade | mesic to wet | bees | butterflies (American Holly Azure) |
| <i>Ilex glabra</i> | Inkberry | white | 4-6' | May-June | sun to partial shade | moist to wet | bees, flies | butterflies (American Holly Azure) |
| <i>Ilex vomitoria</i> | Yaupon | white | 10-12' | March-May | sun to partial shade | dry | bees | butterflies |
| <i>Illicium floridanum</i> | Purple Anise | maroon | 8-9' | April-May | shade to partial shade | moist to wet, acidic | hummingbirds, butterflies, bees | |
| <i>Itea virginica</i> | Virginia Sweetspire | white | 3-5' | May-June | sun to partial shade | moist | butterflies, bees | butterflies (American Holly Azure) |
| <i>Magnolia grandiflora</i> | Southern Magnolia | white | to 60' | May-June | shade | moist | beetles, bees | |
| <i>Magnolia virginiana</i> var. <i>virginiana</i> | Sweet Bay | white | 6-60' | April | sun to partial shade | moist to flooded | bees (honeybees), beetles | butterflies (Eastern Tiger Swallowtail, Spicebush Swallowtail) |
| <i>Morella cerifera</i> | Common Wax Myrtle | white | 10-20' | April | sun to partial shade | moist to wet, salt tolerant | bees | butterflies (Red-banded Hairstreak) |
| <i>Rhododendron alabamense</i> | Alabama Azalea | white | 5-6' | March-April | partial shade | moist | butterflies | |
| <i>Rhododendron atlanticum</i> | Coastal Azalea | pinkish white | 3-6' | April-May | partial shade | moist | butterflies (skippers, swallowtails), hummingbirds | butterflies (Striped Hairstreak, Brown Elfin, Gray Comma) |
| <i>Rhododendron austrinum</i> | Florida Azalea | yellowish-orange | 8-10' | April | partial shade | moist | hummingbirds, butterflies | butterflies |
| <i>Sabal minor</i> | Dwarf Palmetto | white | 4-6' | June-July | sun to partial shade | moist to wet | bees (honeybees, others) | |
| <i>Stewartia malacodendron</i> | Silky Camellia | white | to 20' | May-June | sun to partial shade | moist, acidic | bees, butterflies | |
| <i>Styrax americanus</i> var. <i>americanus</i> | American Snowbell | white | 8-10' | April-June | shade to partial shade | moist | bees | |
| <i>Styrax grandifolius</i> | Bigleaf Snowbell | white | to 20' | April-May | shade to partial shade | moist | bees (bumble bees, others), butterflies | |
| <i>Vaccinium crassifolium</i> | Creeping Blueberry | white | 4-8" | April-May | sun to partial shade | moist | bees, butterflies | butterflies |

| Botanical Name | Common Name | Color | Height | Flower Season | Sun | Soil | Visitation by pollinator | Also a host |
|-----------------------|-------------|-------|--------|---------------|--------------|--------------|--|---|
| <i>Viburnum nudum</i> | Possumhaw | white | 6-8' | April-May | shade to sun | moist to wet | butterflies (Baltimore Checkerspot, Red-spotted Purple, Spring Azure, Appalachian Azure, Striped Hairstreak, White M Hairstreak), bees, beetles, flies | butterflies (Baltimore Checkerspot, Spring Azure) |

Perennial Flowers

| | | | | | | | | |
|--|---------------------------|--------------------------------|--------|-------------|------------------------|----------------------------|--|--|
| <i>Asclepias lanceolata</i> | Red Milkweed | orange-red | 3-5' | June-August | sun to partial shade | moist to wet | bees, moths | butterflies (Monarch, Queen) |
| <i>Bigelovia nudata</i> | Rayless Goldenrod | yellow | 15-24" | Aug-Oct | sun to partial shade | moist to wet | bees | |
| <i>Boltonia caroliniana</i> | Carolina Doll's-daisy | white/pink | 12-40" | Aug-Oct | sun to partial shade | moist | bees | butterflies (Skippers) |
| <i>Conradina canescens</i> | Gray Rosemary | light purple | to 3' | January-May | sun to partial shade | dry, sand | bees (blueberry bee, others) | |
| <i>Chrysogonum virginianum</i> var. <i>australe</i> | Gulf Coast Green-and-gold | yellow | 4-6" | March-May | partial shade | moist to submoist | bees, flies | |
| <i>Coreopsis falcata</i> | Pool Coreopsis | yellow | 25-80" | May-July | sun to partial shade | wet | butterflies (Sulphur) | butterflies |
| <i>Eurybia spectabilis</i> | Showy Aster | purple with yellow center | 1-2' | Aug-Oct | sun to partial shade | dry | butterflies, bees | butterflies |
| <i>Helenium pinnatifidum</i> | Savanna Sneezeweed | yellow | 8-40" | April-May | sun to partial shade | moist to wet | bees, butterflies | |
| <i>Helianthus angustifolius</i> | Narrowleaf Sunflower | yellow | 5-7' | Sept-Oct | sun | moist to wet | butterflies (White Indigo Duskywing, Buckeye, Silvery Checkerspot), bees | butterflies (Silvery Checkerspot, Gorgone Checkerspot, Painted Lady) |
| <i>Helianthus heterophyllus</i> | Savanna Sunflower | yellow | 8-15" | Aug-Oct | sun to partial shade | wet | bees | |
| <i>Hibiscus coccineus</i> | Scarlet Hibiscus | red | to 3' | July-Aug | sun | wet | butterflies (Checked Skipper) | butterflies |
| <i>Hibiscus grandiflorus</i> | Swamp Rose-mallow | pink/white | 4-6' | July-Aug | sun | moist to wet | flies (syrphid flies) | |
| <i>Hibiscus moscheutos</i> | Eastern Rose-mallow | white to pink with red throats | 5-7' | June-Aug | sun | moist to wet | hummingbirds, butterflies (Cloudless Sulphur, Dukes' Skipper) | butterflies (Painted Lady) |
| <i>Iris verna</i> var. <i>verna</i> | Coastal Plain Dwarf Iris | blue with golden yellow sepal | 6-12" | March-May | shade to partial shade | dry | bees | |
| <i>Iris virginica</i> | Southern Blueflag | blue | 24-40" | April-May | sun | wet to moist | butterflies (Sliver Spotted Skipper, Palamedes Swallowtail, Two-spotted Skipper), bumblebees, beetles | |
| <i>Liatris elegans</i> | Elegant Blazing-star | rosy-pink | 3-5' | Sept-Oct | sun to partial shade | dry | butterflies, bees | |
| <i>Lobelia elongata</i> | Blue Lobelia | blue | 24-36" | Aug-Oct | sun to partial shade | mesic to wet, inundated | butterflies | |
| <i>Monarda punctata</i> var. <i>punctata</i> | Eastern Horse-mint | yellow/purplish-pink | 15-24" | Aug-Sept | sun to partial shade | dry, sandy | bees (bumblebees, honeybees, Miner bees, Plasterer bees), moths, butterflies (Silver-spotted Skipper), hummingbirds, wasps | moths (Gray Marvel, Pyralid) |
| <i>Pontederia cordata</i> var. <i>cordata</i> | Pickeralweed | purple | 1-3' | June-Aug | sun to partial shade | wet or inundated | butterflies (skippers) | |
| <i>Pityopsis graminifolia</i> var. <i>tenuifolia</i> | Narrow-leaved Silkgrass | yellow | 8-15" | Sept-Oct | sun to partial shade | dry | bees (bumblebees, sweatbees) | butterflies |
| <i>Rhexia alifanus</i> | Smooth Meadow-beauty | pink | 20-40" | May-Sept | sun to partial shade | moist to wet, inundated | bees | |
| <i>Solidago sempivirens</i> var. <i>mexicana</i> | Seaside Goldenrod | yellow | 1-6' | Sept-Nov | sun | moist to wet | butterflies, beetles, bees | butterflies |
| <i>Stokesia laevis</i> | Stoke's Aster | blue | 1-2' | July-Aug | sun to partial shade | moist | butterflies (Great Spangled Fritillary), honeybees | |
| <i>Yucca filamentosa</i> | Spoonleaf Yucca | white/green | 3-9' | April-June | sun | dry to moist, well drained | moths (yucca moths) | |
| <i>Zephyranthes atamasca</i> | Atamasco Lily | white | 3-12" | March-April | sun to partial shade | moist to wet | moths (hawk moths?) | |

Vines

| | | | | | | | | |
|--------------------------------|--------------------------|-------------------------|--------|-------------|----------------------|--------------|--|---|
| <i>Ampelaster carolinianus</i> | Climbing Aster | rose-pink | 10-20' | Sept-Oct | sun to partial shade | moist to dry | bees (sweatbees, leafcutter bees) | |
| <i>Bignonia capreolata</i> | Cross-vine | red and yellow | 30-45' | April-May | sun to partial shade | moist | hummingbirds, bumblebees | |
| <i>Campsis radicans</i> | Trumpet-creeper | orange | 30-45' | June-July | sun | moist to dry | hummingbirds, bumblebees | moths (Plebeian Sphinx, Pyralid) |
| <i>Clematis crispa</i> | Marsh Clematis | bluish-purple | 10-20' | June-August | sun | moist to wet | bees, butterflies | |
| <i>Wisteria frutescens</i> | American Wisteria | bluish-purple, lavender | to 30' | April-May | sun to partial shade | moist to wet | butterflies (Juvenal's Duskywing) | butterflies (Silver-spotted Skipper, Long-tailed Skipper) |
| <i>Gelsemium rankinii</i> | Swamp Jessamine | yellow | 10-20' | March-April | sun | moist to wet | bees (aphid bee, blueberry bee, others), flies | |
| <i>Ipomoea sagittata</i> | Arrow-leaf Morning Glory | pink | to 15' | July-Sept | sun | moist to wet | bees (bumblebees, others), hummingbirds | |

HABITAT HINTS

FOR THE OUTER COASTAL PLAIN MIXED PROVINCE



HABITAT REQUIREMENTS FOR BEE-POLLINATED GARDEN FLOWERS AND CROPS

| | Bumble | Digger | Lg Carpenter | Sm Carpenter | Squash/Gourd | Leafcutter | Mason | Sweat | Plasterer | Yellow-faced | Andrenid |
|------------------------|--------|--------|--------------|--------------|--------------|------------|-------|-------|-----------|--------------|----------|
| FLOWERS | | | | | | | | | | | |
| Catalpa | | | x | | | | | | | | |
| Catnip | x | x | | | | | x | | | | |
| Clover | | x | | | | | | | | | x |
| Columbine | x | | | | | | | | | | |
| Cow parsley | | | | | | | | | | x | |
| Goldenrod | x | x | | | | x | | x | | | |
| Impatiens | x | | | | | | | | | | |
| Irises | x | | x | | | | | | | | |
| Lavender | x | x | x | | | x | | | | | |
| Milkwort | | | | | | | | x | | | |
| Morning glory | | | | x | | | | | | | |
| Penstemon | x | x | | | | | x | | | | |
| Passion flowers | | | x | | | | | | | | |
| Phacelia | x | x | | x | | x | x | x | x | | x |
| Potentilla | | | | | | | | | | x | |
| Rose | x | | x | | | | x | x | | x | |
| Salvia | x | x | x | | | x | x | | | | |
| Saxifrages | | | | | | | | x | | x | |
| Sorrel | | | | x | | | | | | | |
| Sunflowers | x | x | x | x | | x | | x | x | | x |
| Violet | | | | | | | | x | | | x |
| Wild Mustard | | x | | | | | | | x | | |
| Willow catkins | | | | | | | | | x | | x |
| CROPS | | | | | | | | | | | |
| Almond | x | | | | | | x | | | | x |
| Apple | | | | | | | x | | | | |
| Blueberry | x | x | | | | | | | | | x |
| Cherry | | | | | | | x | | | | x |
| Eggplant | x | | x | | | | | x | | | |
| Gooseberry | x | | | | | | | | | | x |
| Legumes | x | x | | | | x | | x | | | |
| Water melon | x | | | | | | | x | | | |
| Squash/Pumpkins/Gourds | | | x | | x | | | | | | |
| Tomatoes | x | x | x | | | | | x | | | |
| Thyme | x | x | | | | | x | x | | x | |



HABITAT AND NESTING REQUIREMENTS:

Bumble Bees:

Abandoned mouse nests, other rodent burrows, upside down flower pots, under boards, and other human-made cavities. Colonies are founded by a queen in the spring and don't die out in the fall. New queens mate then and overwinter in a sort of hibernation. Bumble bees are usually active during the morning hours and forage at colder temperatures than honey bees, even flying in light rain.

Large carpenter bees:

Soft dead wood, poplar, cottonwood or willow trunks and limbs, structural timbers including redwood. Depending on the species, there may be one or two brood cycles per year. These bees can be active all day even in the hottest weather.

Digger bees:

Sandy soil, compacted soils, bank sides. Anthophorid bees (now in the Apidae) are usually active in the morning hours, but can be seen at other times.

Small carpenter bees:

Pithy stems including roses and blackberry canes. These bees are more active in the morning but can be found at other times.

Squash and Gourd bees:

Sandy soil, may nest in gardens (where pumpkins, squash and gourds are grown) or pathways. These bees are early risers and can be found in pumpkin patches before dawn. Males often sleep in the wilted flowers.

Leafcutter bees:

Pre-existing circular tunnels of various diameters in dead but sound wood created by emerging beetles, some nest in the ground. Leave dead limbs and trees to support not just pollinators but other wildlife. Leafcutter bees can be seen foraging throughout the day even in hot weather.

Mason bees:

Pre-existing tunnels, various diameters in dead wood made by emerging beetles, or human-made nesting substrates, drilled wood boards, paper soda straws inserted into cans attached to buildings. Mason bees are generally more active in the morning hours.

Sweat bees:

Bare ground, compacted soil, sunny areas not covered by vegetation. Like most bees, sweat bees forage for pollen earlier in the morning and then for nectar later.

Plasterer or cellophane bees:

Bare ground, banks or cliffs. Colletid bees can be active in the morning or later in the day.

Yellow-faced bees:

In dead stems. These bees are more active during morning hours.

Andrenid bees:

Sunny, bare ground, sand soil, under leaf litter or in soil in banksides and cliffs. These generally spring-active bees are most commonly seen on flowers during the morning when pollen and nectar resources are abundant.



**“MONARCH
BUTTERFLIES
NEVER FAIL TO
CATCH THE
VISITOR’S EYE
AND ALWAYS
LEAD TO
A TEACHABLE
MOMENT.”**

-- LOGAN LEE,
PRAIRIE SUPERVISOR
MIDWIN NATIONAL
TALLGRASS PRAIRIE

BECOME FAMILIAR WITH POLLINATORS IN YOUR LANDSCAPE.

- ✿ Watch for activity throughout the day and the seasons.
- ✿ Keep a simple notebook of when and what comes to your garden.
NOTE: It is not necessary to identify each species when you first get started. Simply note if it is a bee that likes the yellow flower that blooms in the fall.
- ✿ Consult a local field guide or web site when you are ready to learn more details.

ADD NATIVE PLANTS TO ATTRACT MORE NATIVE POLLINATORS.

- ✿ List the plants you currently have in your landscape.
- ✿ Determine when you need additional flowers to provide nectar and pollen throughout the growing season.
- ✿ Add plants that provide additional seasons of bloom, create variable heights for shelter, and attract the types of pollinators you want.
- ✿ Don't forget to include host plants that provide food and shelter for larval development.
- ✿ Contact your local native plant society or extension agent for more help.

USE POLLINATOR FRIENDLY LANDSCAPE PRACTICES TO SUPPORT THE POLLINATORS YOU ATTRACT.

- ✿ Use Integrated Pest Management Practices to address pest concerns.
- ✿ Tolerate a little mess – leave dead snags and leaf litter, keep areas bare for ground nesting insects, and leave some weeds that provide food for pollinators.
- ✿ Provide safe access to clean water.

NOTICE THE CHANGES THAT YOU HAVE HELPED TO CREATE!



RESOURCES

Many books, websites, and people were consulted to gather information for this guide. Use this list as a starting point to learn more about pollinators and plants in your area.

BAILEY'S ECOREGION MAPS

USDA Forest Service
fs.fed.us/land/ecosysmgmt/ecoreg1_home.html

POLLINATION/POLLINATORS

Pollinator Partnership
pollinator.org

North American Pollinator Protection Campaign
nappc.org

Bee Friendly Gardening
pollinator.org/bfg

Natural Resources Conservation Service
nrcs.usda.gov

USDA Forest Service
fs.fed.us/wildflowers/pollinators/

Wild Farm Alliance
wildfarmalliance.org

Xerces Society Pollinator Program
xerces.org

Illinois Natural History Survey
inhs.uiuc.edu

The Forgotten Pollinators
Buchmann, S.L. and G.P. Nabhan. 1997. Island Press: Washington, DC.

Status of Pollinators in North America
2007. The National Academies Press: Washington, DC.

NATIVE PLANTS/GARDENING

Plant Conservation Alliance
nps.gov/plants

Seeds of Success
nps.gov/plants/sos

Lady Bird Johnson Wildflower Center
wildflower.org/plants/

USDA Hardiness Zone Map
usna.usda/Hardzone/

U.S. National Arboretum
usna.usda.gov/Hardzone/ushzmap.html

USDA, NRCS. 2007. The PLANTS Database
plants.usda.gov

NATIVE BEES

Alternative Pollinators: Native Bees National Sustainable Information Service
by Lane Greer, NCAT Agriculture Specialist, Published 1999, ATTRA Publication #IP126
attra.ncat.org/attra-pub/nativebee.html

Plants Attractive to Native Bees Agriculture Research Service
ars.usda.gov/Research/docs.htm?docid=12052

Bees of the World
Christopher O'Toole and Anthony Raw. 1999. Blandford. London, UK.

BUTTERFLIES AND MOTHS

Butterflies and Moths of North America.

Opler, Paul A., Harry Pavulaan, Ray E. Stanford, Michael Pogue, coordinators. 2006.
butterfliesandmoths.org/

Butterflies of North America
Jim Brock and Kenn Kaufman. 2003. Houghton Mifflin. New York, NY.

North American Butterfly Association
naba.org

FEEDBACK

We need your help to create better guides for other parts of North America. Please e-mail your input to feedback@pollinator.org or fax to 415-362-3070.

- ✿ How will you use this guide?
- ✿ Do you find the directions clear? If not, please tell us what is unclear.
- ✿ Is there any information you feel is missing from the guide?
- ✿ Any other comments?

THANK YOU FOR TAKING THE TIME TO HELP!

POLLINATOR PARTNERSHIP

Protect their lives. Preserve ours.



NAPPC



Plant Conservation Alliance



RESEARCH AND WRITING:

ELIZABETH L. LEY
STEPHEN BUCHMANN, PH.D.
GARY KAUFFMAN
KATHERINE MCGUIRE

EDITORIAL:

LAURIE DAVIES ADAMS
AND LARRY STRITCH, PH.D.

PRODUCTION SUPERVISION:

KATHERINE MCGUIRE

DESIGN:

MARGUERITE MEYER

CONCEPT REVIEW:

American Farm Bureau Federation, Ron Gaskell
Bureau of Land Management, Peggy Olwell, Carol Spurrier,
Mary Byrne, Mary Tisdale, Elizabeth Wooster
National Garden Association, Susanne DeJohn
Plant Conservation Alliance – Edward Fletcher, Jean Giblette,
Mary Ann Lawler, Ron Smith
Smithsonian Institute, Department of Botany,
Gary Krupnick, Ph.D.
USDA - CSREES, Greg Crosby, Ph.D., Leslie Gilbert, Ph.D.
USDA - Forest Service, David Pivorunas, Larry Stritch, Ph.D.
USDA - Natural Resource Conservation Service, Doug Holy,
Hilda Diaz-Soltero
USDOI - US Fish and Wildlife Service, Karen Anderson,
Don MacLean, Patricia DeAngelis, Ph.D.
USGS - Steve Hilburger, Elizabeth Sellers

PHOTO CONTRIBUTOR:

dangphoto.net

ILLUSTRATIONS:

Carolyn Vibbert

For a copy of this brochure, or for another region, visit pollinator.org

Pollinator Partnership/North American Pollinator Protection Campaign

pollinator.org  nappc.org  Email: info@pollinator.org

